

STABLE REDUCTION OF FINITE COVERS OF CURVES

QING LIU

ABSTRACT. Let K be the function field of a connected regular scheme S of dimension 1, and let $f : X \rightarrow Y$ be a finite cover of projective smooth and geometrically connected curves over K with $g(X) \geq 2$. Suppose that f can be extended to a finite cover $\mathcal{X} \rightarrow \mathcal{Y}$ of semi-stable models over S (it is known that this is always possible up to finite separable extension of K). Then there exists a unique minimal such cover. This gives a canonical way to extend $X \rightarrow Y$ to a finite cover of semi-stable models over S .

Let S be a Dedekind scheme (i.e a connected Noetherian regular scheme of dimension 1), with field of functions $K := K(S)$. Let $f : X \rightarrow Y$ be a finite morphism of smooth geometrically connected projective curves over K . We can ask how to extend, in some canonical way, the morphism f to a morphism of models of X and Y over S . It is proved in [15], 4.4 that if X and Y have stable models $\mathcal{X}^{\text{st}}, \mathcal{Y}^{\text{st}}$ over S , then f extends uniquely to a morphism $\mathcal{X}^{\text{st}} \rightarrow \mathcal{Y}^{\text{st}}$. However we will in general lose the finiteness of f . On the other hand, after a finite separable extension of K , f extends to a finite morphism of semi-stable models $\mathcal{X} \rightarrow \mathcal{Y}$ over S ([4]; [15], Remark 4.6; and Corollary 3.10 below). Following Coleman [4], such a pair $\mathcal{X} \rightarrow \mathcal{Y}$ is called a *semi-stable model* of f , and it is called *stable* if moreover it is minimal among the semi-stable models of f (cf. 3.1). The stable model of f (if it exists) is unique up to isomorphism.

Theorem 0.1. (Corollary 4.6) *Suppose that either $g(X) \geq 2$, or $g(X) = 1$ and X has potentially good reduction. Then there exists a finite separable extension K' of K such that $X_{K'} \rightarrow Y_{K'}$ admits a stable model $\mathcal{X}' \rightarrow \mathcal{Y}'$ over S' , where S' is the integral closure of S in K' . Moreover, for any Dedekind scheme T dominating S' , $\mathcal{X}'_T \rightarrow \mathcal{Y}'_T$ is the stable model of $X_{K(T)} \rightarrow Y_{K(T)}$ over T .*

This gives a canonical way to extend a finite cover of projective smooth curves over K to a finite cover of semi-stable models over (some finite cover of) S . The last part of the theorem says that the stable model of f commutes with flat base change. The stable model of f should be seen as an analogue of the stable model of a curve. In a forthcoming work, we will apply this theorem to study a compactification of Hurwitz moduli spaces of finite covers of curves. We also prove the theorem for smooth marked curves X, Y . Note that in general $\mathcal{X}', \mathcal{Y}'$ are not the respective stable models of the curves $X_{K'}, Y_{K'}$. The proof of 0.1 is based on a more general result:

Theorem 0.2. (Theorem 4.5) *Let $f : X \rightarrow Y$ be a finite morphism of smooth geometrically connected projective curves over K . Let $\mathcal{X}, \mathcal{Y}$ be respective models of X and Y over S . Then there exists a finite separable extension K' of K , and models $\mathcal{X}', \mathcal{Y}'$ of $X_{K'}$ and $Y_{K'}$ over S' (integral closure of S in K') such that*

2000 *Mathematics Subject Classification.* 14G20, 14H30, 14D10, 11G20.

Key words and phrases. curves, finite covers, stable reduction.

- (1) $\mathcal{X}', \mathcal{Y}'$ dominate respectively $\mathcal{X}_{S'}, \mathcal{Y}_{S'}$ and are semi-stable over S' ;
- (2) $X_{K'} \rightarrow Y_{K'}$ extends to a finite morphism $\mathcal{X}' \rightarrow \mathcal{Y}'$;
- (3) the pair $(\mathcal{X}', \mathcal{Y}')$ is minimal : for any pair $(\mathcal{X}'', \mathcal{Y}'')$ satisfying properties (1) and (2), $\mathcal{X}''$ and $\mathcal{Y}''$ dominate respectively $\mathcal{X}'$ and $\mathcal{Y}'$. The cover $\mathcal{X}' \rightarrow \mathcal{Y}'$ is called the stable hull of $\mathcal{X}_{S'} \dashrightarrow \mathcal{Y}_{S'}$.
- (4) the formation of $\mathcal{X}', \mathcal{Y}'$ commutes with flat base change $T \rightarrow S'$: if T is a Dedekind scheme dominating S' , then $\mathcal{X}'_T \rightarrow \mathcal{Y}'_T$ is the stable hull of $\mathcal{X}_T \dashrightarrow \mathcal{Y}_T$.

The paper is organized as follows. In Section 1, $S = \text{Spec } \mathcal{O}_K$ is local. We discuss some sufficient conditions for a model of $X_{\hat{K}}$ over $\hat{\mathcal{O}}_K$ (completion of $\mathcal{O}_K$) to be defined over $\mathcal{O}_K$. E.g., this is true for semi-stable or regular models. This is used to reduce the proof of 4.4 to the case of a complete local base S . In Section 2, we prove the special case of 0.2 when $X = Y, \mathcal{X} = \mathcal{Y}$ and $f = \text{Id}$. Parts (1) and (2) of 0.2 are proved in Section 3. The existence of a stable model (Part (3)) is proved in Section 4.

Convention Through this paper, S is a Noetherian regular connected scheme of dimension 1 (Dedekind scheme), and, unless otherwise specified, X, Y are projective, smooth and geometrically connected curves over $K := K(S)$. When we state that a property (P) holds for some model $\mathcal{X}$ of X after finite separable extension of K , this means that there exists a finite separable extension of K'/K , such that (P) is satisfied for $\mathcal{X}_{S'}$, where S' is the normalization of S in K' .

Note that the hypothesis X, Y geometrically connected (instead of connected) is not serious. Actually X is geometrically connected over the finite separable extension $H^0(X, \mathcal{O}_X)$ of K .

ACKNOWLEDGEMENTS

I would like to thank D. Lorenzini for pointing out Coleman's paper [4] and for interesting comments, Ch. Deninger and A. Werner for motivational discussions and Sylvain Maugeais for pointing out some mistakes in an earlier version. I also thank the referee to correct a mistake in the proof of Proposition 4.11.

1. DESCENT FROM THE COMPLETION

Let $\mathcal{O}_K$ be a discrete valuation ring, and let X be a geometrically connected smooth projective curve K . We give some sufficient conditions for a model of $X_{\hat{K}}$ over $\hat{\mathcal{O}}_K$ to be defined over $\mathcal{O}_K$.

Lemma 1.1. *Let $\mathcal{O}_K$ be a discrete valuation ring, let A be a flat local $\mathcal{O}_K$ -algebra, localization of a finitely generated $\mathcal{O}_K$ -algebra. Suppose that $\text{Spec}(A \otimes \hat{K})$ is integral, one-dimensional, and smooth over $\hat{K}$. Let $\mathcal{W} \rightarrow \text{Spec } \hat{A}$ be a projective birational morphism. Then under any of the following conditions, $\mathcal{W}$ and the morphism $\mathcal{W} \rightarrow \text{Spec } \hat{A}$ are defined over A :*

- (a) $\text{Pic}(\hat{A} \otimes \hat{K})$ is a torsion group (i.e., every element has finite order);
- (b) $\hat{A}$ is $\mathbb{Q}$ -factorial.

Proof: The morphism $\mathcal{W} \rightarrow \text{Spec } (\hat{A})$ is the blowing-up along a closed subscheme $V(\mathfrak{J})$ of $\text{Spec } (\hat{A})$. Let t be a uniformizing element of $\mathcal{O}_K$. Let us show that $\mathfrak{J}$ can be chosen in such a way that $t^n \in \mathfrak{J}$ for some $n \geq 1$. Let us suppose that condition (a) is satisfied. Then there exists some $m \geq 1$ such that the restriction of $\mathfrak{J}^m$ to

the generic fiber is principal, generated by a $f \in \mathfrak{I}^m \otimes \hat{K}$. Under condition (b), let D be the scheme theoretical closure of $V(\mathfrak{I}) \cap \text{Spec}(\hat{A} \otimes \hat{K})$ in $\text{Spec}(\hat{A})$, considered as a Weil divisor. Then mD is principal for some $m > 0$, generated by a $f \in \mathfrak{I}^m$.


In both cases, there exist $a, b \in \mathbb{Z}$ such that $t^a \mathfrak{I}^m \subseteq f \hat{A} \subseteq t^b \mathfrak{I}^m$. Replacing $\mathfrak{I}$ by the ideal $t^a f^{-1} \mathfrak{I}^m$ (which does not change the blowing-up along $\mathfrak{I}$), we can suppose that the restriction of $\mathfrak{I}$ to the generic fiber is trivial, and $t^n \in \mathfrak{I}$ for some $n \geq 0$ ($n = a - b$). Since $\mathcal{O}_K$ is dense in $\hat{\mathcal{O}}_K$, $\mathfrak{I}$ is then generated by an ideal I of A . Let $\mathcal{X} \rightarrow \text{Spec } A$ be the blowing-up along $V(I)$, then $\mathcal{W} \rightarrow \text{Spec}(\hat{A})$ is obtained from $\mathcal{X} \rightarrow \text{Spec } A$ by the base change $\hat{\mathcal{O}}_K/\mathcal{O}_K$. $\square$

Remark 1.2 Let $\mathcal{W}$ be a projective regular model of $X_{\hat{K}}$ over $\hat{\mathcal{O}}_K$ dominating $\mathcal{X}_{\hat{\mathcal{O}}_K}$ for some model $\mathcal{X}$ of X over $\mathcal{O}_K$. Then $\mathcal{W}$ and $\mathcal{W} \rightarrow \mathcal{X}_{\hat{\mathcal{O}}_K}$ are defined over $\mathcal{O}_K$. Actually, if $\tilde{\mathcal{X}} \rightarrow \mathcal{X}$ is the minimal desingularization of $\mathcal{X}$, then $\tilde{\mathcal{X}}_{\hat{\mathcal{O}}_K} \rightarrow \mathcal{X}_{\hat{\mathcal{O}}_K}$ is the minimal desingularization of $\mathcal{X}_{\hat{\mathcal{O}}_K}$ (same proof as in [13], 9.3.28). Hence $\mathcal{W} \dashrightarrow \tilde{\mathcal{X}}_{\hat{\mathcal{O}}_K}$ is a birational morphism. It is a sequence of blowing-ups of closed points ([13], 9.2.2). Hence $\mathcal{W}$ is defined over $\mathcal{O}_K$, and so is the morphism $\mathcal{W} \rightarrow \mathcal{X}_{\hat{\mathcal{O}}_K}$ by faithfully flat descent of rational maps.

Proposition 1.3. *Let $S = \text{Spec } \mathcal{O}_K$ be local. Let $\mathcal{X}$ be a model of X over S dominating some semi-stable or regular model. Let $\varphi : \mathcal{W} \rightarrow \mathcal{X}_{\hat{\mathcal{O}}_K}$ be a projective birational morphism over $\hat{\mathcal{O}}_K$. Then $\mathcal{W}$ and φ are defined over S .*

Proof: It is enough to show that $\mathcal{W}$ is defined over S . We can suppose that $\mathcal{X}$ itself is semi-stable or regular. The morphism φ is an isomorphism outside of a finite set F of closed points of $(\mathcal{X}_{\hat{\mathcal{O}}_K})_s = \mathcal{X}_s$. Let $x \in F$, and let $A_x = \mathcal{O}_{\mathcal{X},x}$. If $\mathcal{X}$ is regular, then $\hat{A}_x$ is regular and thus factorial. If $\mathcal{X}$ is semi-stable, there exists a finite (étale) extension $\mathcal{O}_L/\hat{\mathcal{O}}_K$ such that $\hat{A}_x \otimes_{\hat{\mathcal{O}}_K} \mathcal{O}_L$ is a finite direct sum of rings $\mathcal{O}_L[[u]]$ or $\mathcal{O}_L[[u, v]]/(uv - a)$, $a \in \mathcal{O}_L$. It is well-known that Pic of these rings tensored by $\hat{L}$ are trivial (see for instance [9], Cor. 2.2, for $\mathcal{O}_L[[u, v]] \otimes \hat{L}/(uv - a)$). Hence $\text{Pic}(\hat{A}_x)$ is torsion ([13], 7.2.18, 7.2.19). By Lemma 1.1, $\mathcal{W} \times_{\mathcal{X}_{\hat{\mathcal{O}}_K}} \text{Spec } \hat{A}_x \rightarrow \text{Spec } \hat{A}_x$ is defined over A_x . By glueing the morphisms above $\text{Spec } \hat{A}_x$, when x varies in F , and the isomorphism above $\mathcal{X} \setminus F$, we find a morphism over $\mathcal{X}$ which is equal to φ when base changed to $\hat{\mathcal{O}}_K$. $\square$

Remark 1.4 In general, not every birational projective morphism $\mathcal{W} \rightarrow \mathcal{X}_{\hat{\mathcal{O}}_K}$ is defined over $\mathcal{O}_K$. In other words, even if $\mathcal{W} \rightarrow \mathcal{X}_{\hat{\mathcal{O}}_K}$ is an isomorphism outside of the special fiber, it is not necessarily the blowing-up along a closed subscheme with support in the special fiber. To construct such a counterexample, we will imitate the example of a non contractible component given in [2], Lemma 6.7.6. Let us consider the smooth elliptic curve $\mathcal{E}/\mathcal{O}_K$ and the point $a_k \in \mathcal{E}_k(k)$ as given in [2], p. 171. The point a_k satisfies the property that no multiple (in the sense of the group law on $\mathcal{E}$) na_k , $n > 0$, can be lifted to a section in $\mathcal{E}(\mathcal{O}_K)$.



Let $\mathcal{X}' \rightarrow \mathcal{E}$ be the blowing-up of $\mathcal{E}$ along $\{a_k, e_k\}$, where e is the unit section of $\mathcal{E}$. Let P_a and P_e be the respective inverse images of a_k, e_k in $\mathcal{X}'$. They are projective lines over k . Let $\tilde{\mathcal{E}}_k$ be the strict transform of $\mathcal{E}_k$ in $\mathcal{X}'$. The unit section of E_K gives rise to a section of $\mathcal{X}'$ which meets $\mathcal{X}'_k$ at an interior point of P_e . Hence there exists a contraction map $\mathcal{X}' \rightarrow \mathcal{X}$ of $P_a \cup \tilde{\mathcal{E}}_k$ ([2], Corollary 6.7.3). Now over $\hat{\mathcal{O}}_K$, we can contract $\tilde{\mathcal{E}}_k$ in $\mathcal{X}'_{\hat{\mathcal{O}}_K}$ (op. cit., 6.7.4), which gives a model $\mathcal{W}/\hat{\mathcal{O}}_K$ and we have birational projective morphisms $\mathcal{X}'_{\hat{\mathcal{O}}_K} \rightarrow \mathcal{W} \rightarrow \mathcal{X}_{\hat{\mathcal{O}}_K}$. We want to show that $\mathcal{W}$ is not defined over $\mathcal{O}_K$, or, equivalently, that $\tilde{\mathcal{E}}_k$ can not be contracted over $\mathcal{O}_K$. Suppose that $\mathcal{W}$ exists over $\mathcal{O}_K$, then there exists a relative effective Cartier divisor D on $\mathcal{X}'$ such that $\text{Supp} D$ meets P_a and P_e but not $\tilde{\mathcal{E}}_k$. Taking the Zariski closure of D_K in $\mathcal{E}$ gives a relative effective Cartier divisor D' on $\mathcal{E}$ such that $D'_k = na_k + me_k$ for some integers $n, m > 0$. But $D' - (n+m)e \in \text{Pic}^0_{\mathcal{E}/\mathcal{O}_K}(\mathcal{O}_K) \simeq \mathcal{E}(\mathcal{O}_K)$ then defines a section $b \in \mathcal{E}(\mathcal{O}_K)$ such that $D' - (n+m)e \sim b - e$. In the special fiber we then have $n(a_k - e_k) \sim b_k - e_k$, hence $na_k = b_k$ in the group $\mathcal{E}_k(k)$, contradiction with the assumption on a_k .

Lemma 1.5. *Suppose that X has semi-stable reduction over S . Then every relatively minimal semi-stable model $\mathcal{W}$ of $X_{\hat{K}}$ over $\hat{\mathcal{O}}_K$ is defined over $\mathcal{O}_K$.*

Proof: If $g(X) \geq 2$, then the unique relatively minimal semi-stable model of X is the stable model of X over S . Since the stable model is unique and commutes with flat base change ([13], 10.3.36), $\mathcal{W}$ over $\hat{\mathcal{O}}_K$ is defined over $\mathcal{O}_K$. The same is true if $g(X) = 1$ and X has good reduction.

Suppose that $g(X) = 1$ and X has multiplicative reduction. Let $\mathcal{X}$ be a relatively minimal semi-stable model. Let $\tilde{\mathcal{X}}$ be its minimal desingularization. Let us show that $\tilde{\mathcal{X}}$ is the minimal regular model of X over S . Let Γ be an irreducible component of $\mathcal{X}_s$ and let $\tilde{\Gamma}$ be its strict transform in $\tilde{\mathcal{X}}$. By Lemma 2.13(a), $\deg \omega_{\mathcal{X}/S}|_{\Gamma} > 0$ if $\mathcal{X}_s$ is reducible. If $\mathcal{X}_s$ is irreducible, then $\deg \omega_{\mathcal{X}/S}|_{\Gamma} = 2g(X) - 2 = 0$. So in any case $\deg \omega_{\tilde{\mathcal{X}}/S}|_{\tilde{\Gamma}} = \deg \omega_{\mathcal{X}/S}|_{\Gamma} \geq 0$. Hence $\tilde{\mathcal{X}}$ has no exceptional divisor. Since any strict subset of the set of irreducible components of $\tilde{\mathcal{X}}_s$ can be contracted ([13], 9.4.19) into a semi-stable model (2.13(a)), $\mathcal{X}_s$ is irreducible. So the relatively minimal semi-stable models of X correspond bijectively to the irreducible components of the minimal regular model of X . The same is true over $\hat{\mathcal{O}}_K$. Since the minimal regular model commutes with the base change $\hat{\mathcal{O}}_K/\mathcal{O}_K$ ([13], 9.3.28), we see that the relatively minimal semi-stable models of $X_{\hat{K}}$ are exactly those of X base changed to $\hat{\mathcal{O}}_K$.

Suppose that $g(X) = 0$. Let $\mathcal{X}$ be a relatively minimal semi-stable model of X over S and let $\Gamma_1, \dots, \Gamma_n$ be the irreducible components of $\mathcal{X}_s$. Then

$$\sum_i \deg \omega_{\mathcal{X}/S}|_{\Gamma_i} = 2g(X) - 2 < 0.$$

So $\omega_{\mathcal{X}/S}$ has negative degree on at least one component Γ_i . By Lemma 2.13(a), $n = 1$, thus $\mathcal{X}_s$ is irreducible, semi-stable and has arithmetic genus 0. So $\mathcal{X}_s$ is smooth. The same reasoning shows that $\mathcal{W}$ is smooth. Let $\mathcal{L}$ be the dual of the dualizing sheaf on $\mathcal{W}$. Let $s_0, s_1 \in H^0(\mathcal{W}, \mathcal{L})$ be a basis with $s_i \in H^0(X, \omega_{X/K}^\vee)$. Then s_0, s_1 define a closed immersion $\mathcal{W} \rightarrow \mathbb{P}^2$ over $\hat{\mathcal{O}}_K$. Its image is a conic defined by a polynomial with coefficients in K . Hence $\mathcal{W}$ is defined over $\mathcal{O}_K$. $\square$

The next corollary is weaker than 1.3, but sufficient for the purpose of Theorem 4.5. It is an immediate consequence of 1.3 and the above lemma. However we give a direct proof without using 1.3.

Corollary 1.6. *Let $S = \text{Spec } \mathcal{O}_K$ be local. Let X be a smooth geometrically connected projective curve over K . Then every semi-stable model $\mathcal{W}$ of $X_{\hat{K}}$ over $\hat{\mathcal{O}}_K$ is defined over $\mathcal{O}_K$.*

Proof: The model $\mathcal{W}$ dominates a relatively minimal semi-stable model $\mathcal{Z}$ of $X_{\hat{K}}$ which is then defined over $\mathcal{O}_K$ by 1.5. Let $\widetilde{\mathcal{W}} \rightarrow \mathcal{W}$ be the minimal desingularization of $\mathcal{W}$. Then $\widetilde{\mathcal{W}} \rightarrow \mathcal{Z}$ is defined over $\mathcal{O}_K$ (1.2). The irreducible components of $\widetilde{\mathcal{W}}_s$ in the exceptional locus of $\widetilde{\mathcal{W}} \rightarrow \mathcal{W}$ are (-2) -curves and can be contracted over $\mathcal{O}_K$ (2.13(a)). Hence $\mathcal{W}$ is defined over $\mathcal{O}_K$. $\square$

Remark 1.7 If $X = \mathbb{P}_K^1$, then every normal model $\mathcal{W}$ of $X_{\hat{K}}$ is defined over $\mathcal{O}_K$. Indeed, if $\widetilde{\mathcal{W}}$ is the minimal desingularization of $\mathcal{W}$, then $\widetilde{\mathcal{W}}$ dominates a relatively minimal regular model. The latter is smooth and defined over $\mathcal{O}_K$. Hence $\widetilde{\mathcal{W}}$ is defined over $\mathcal{O}_K$. Now every strict subset of the set of irreducible components of $\widetilde{\mathcal{W}}_s$ is contractible over $\mathcal{O}_K$ ([13], Exerc. 9.4.5). So $\mathcal{W}$ is defined over $\mathcal{O}_K$.

2. STABLE HULL OF A MODEL

Definition 2.1 Let S be a connected Noetherian regular scheme of dimension 1 (i.e., a *Dedekind scheme*). Let X be an integral projective variety over K . A *model* $\mathcal{X}$ of X over S is an integral projective scheme over S whose generic fiber is isomorphic to X . Recall that $\mathcal{X}$ is said to be *semi-stable* if its geometric fibers are reduced with only ordinary double points as singularities. A morphism of models is defined in an obvious way.

Definition 2.2 Let X be a connected projective smooth curve over K , and let $\mathcal{X}$ be a model of X over S . A *stable hull* of $\mathcal{X}$ is a semi-stable model $\mathcal{W}$ of X dominating $\mathcal{X}$, and minimal for these properties (i.e., every semi-stable model dominating $\mathcal{X}$ dominates $\mathcal{W}$).

The aim of this section is to prove the next result.

Theorem 2.3. *Let X be a geometrically connected projective smooth curve over K and let $\mathcal{X}$ be a model of X over S .*

- (a) *The stable hull of $\mathcal{X}$ is unique (up to isomorphism) when it exists. In general, there exists a finite separable extension K'/K such that $\mathcal{X}_{S'}$ (where S' is the integral closure of S in K') has a stable hull over S' .*
- (b) *The stable hull commutes with flat base change: suppose that $\mathcal{X}$ admits a stable hull $\mathcal{W}$ over S and let $S' \rightarrow S$ be a flat morphism of Dedekind schemes, then $\mathcal{W}_{S'}$ is the stable hull of $\mathcal{X}_{S'}$ over S' .*

The proof of the theorem is postponed to 2.17.

Lemma 2.4. *Let G be a finite group acting on X . Let $\mathcal{X}$ be a model of X over S . Then there exists a model $\mathcal{Z}$ of X dominating $\mathcal{X}$, endowed with an action of G , and minimal for these properties.*

Proof: (See [10], 7.6) Let $\sigma \in G$, then there exists a model $\mathcal{X}^\sigma$ such that $\sigma : X \rightarrow X$ extends to an isomorphism $\sigma : \mathcal{X} \rightarrow \mathcal{X}^\sigma$. If $\tau \in G$, by composing $(\sigma^{-1}\tau) : \mathcal{X}^{\tau^{-1}\sigma} \rightarrow \mathcal{X}$ with $\sigma : \mathcal{X} \rightarrow \mathcal{X}^\sigma$, we obtain an isomorphism $\mathcal{X}^{\tau^{-1}\sigma} \rightarrow \mathcal{X}^\sigma$ denoted by τ . Let $\mathcal{P}$ be the fiber product $\prod_{S, \sigma \in G} \mathcal{X}^\sigma$ over S . Then we can make G act on $\mathcal{P}$ by

$$\tau : (x_\sigma)_\sigma \mapsto (\tau(x_{\tau^{-1}\sigma}))_\sigma.$$

Moreover, the diagonal morphism $\Delta : X \rightarrow \mathcal{P}_K, x \mapsto (x, \dots, x)$ is G -equivariant. Let $\mathcal{Z}$ be the Zariski closure of $\Delta(X)$ in $\mathcal{P}$, endowed with the reduced structure. Then G leaves stable $\mathcal{Z}$. Note that $\mathcal{Z}$ dominates $\mathcal{X}$ because the projection morphism $\mathcal{P} \rightarrow \mathcal{X}$ induces a morphism $\mathcal{Z} \rightarrow \mathcal{X}$ which is an isomorphism on the generic fibers.

Let us prove that $\mathcal{Z}$ is minimal. Let $\mathcal{W}$ be a model endowed with an action of G and a birational morphism $\mathcal{W} \rightarrow \mathcal{X}$. Then we have a morphism $\mathcal{W} \rightarrow \mathcal{X}^\sigma$ for all σ and hence a morphism $h : \mathcal{W} \rightarrow \mathcal{P}$. Since $h(\mathcal{W})$ is irreducible and contains $\mathcal{Z}$, h induces a morphism of models $\mathcal{W} \rightarrow \mathcal{Z}$. $\square$

Let us give a corollary of 2.3.

Corollary 2.5. *Let G be a finite group acting on X . Let $\mathcal{X}$ be a model of X over S . Then after finite separable extension of K , $\mathcal{X}$ is dominated by a semi-stable (resp. semi-stable and regular) model $\mathcal{W}$ such that the action of G extends to $\mathcal{W}$. Moreover, there exists a minimal such a model $\mathcal{W}$.*

Proof: Let $\mathcal{Z}$ be the model defined in 2.4. Let $\mathcal{W}$ be the stable hull of $\mathcal{Z}_{S'}$ over some S'/S (Theorem 2.3(a)). By the uniqueness property, the action of G on $\mathcal{Z}_{S'}$ extends to $\mathcal{W}$. It is clear that $\mathcal{W}$ is minimal with respect to the required properties. To have a minimal semi-stable regular model, it is enough to take the minimal desingularization of $\mathcal{W}$. $\square$

Lemma 2.6. *Let S be local with separably closed residue field. Let $\mathcal{X}$ be a model of X with minimal desingularization $\tilde{\mathcal{X}} \rightarrow \mathcal{X}$. Suppose that $\tilde{\mathcal{X}}$ dominates a regular model $\mathcal{Z}$ and that $\mathcal{X}_s, \mathcal{Z}_s$ are geometrically reduced. Then $\tilde{\mathcal{X}}_s$ is geometrically reduced.*

Proof: Note that if $\mathcal{W}$ is a normal model of X , then $\mathcal{W}_s$ verifies the property (S₁), thus $\mathcal{W}_s$ is geometrically reduced if and only if every irreducible component of $\mathcal{W}_s$ has geometric multiplicity ([2], 9.1.3) equal to 1 in $\mathcal{W}_s$. The latter condition depends only on the generic points of $\mathcal{W}_s$. We can decompose $\tilde{\mathcal{X}} \rightarrow \mathcal{Z}$ into a sequence of blowing-ups

$$\tilde{\mathcal{X}} =: \mathcal{Z}_0 \rightarrow \mathcal{Z}_1 \rightarrow \dots \rightarrow \mathcal{Z}_n := \mathcal{Z}$$

such that $\mathcal{Z}_i \rightarrow \mathcal{Z}_{i+1}$ consists in blowing-down an exceptional divisor Θ_i contained in $\mathcal{Z}_i$. We will show by induction that Θ_i has multiplicity 1 in $(\mathcal{Z}_i)_s$ and $\Theta_i \simeq \mathbb{P}_{k(s)}^1$. Since $(\mathcal{Z}_n)_s$ is geometrically reduced, this will imply that $\tilde{\mathcal{X}}_s$ is geometrically reduced.

By the minimality of $\tilde{\mathcal{X}} \rightarrow \mathcal{X}$, Θ_0 is not mapped to a closed point in $\mathcal{X}$. Thus $\tilde{\mathcal{X}} \rightarrow \mathcal{X}$ is an isomorphism in a neighborhood of the generic point of Θ_0 . In particular, Θ_0 has geometric multiplicity 1. Since Θ_0 is a projective line by Castelnuovo criterion, it is isomorphic to $\mathbb{P}_{k(s)}^1$. Suppose that the same holds for $\Theta_j, 0 \leq j \leq i-1$.

Let $\Theta_{i,j}$ be the strict transform of Θ_i in $\mathcal{Z}_j$, $0 \leq j \leq i-1$. If $\Theta_{i,j}$ meets Θ_j for some j , then the computation of $\Theta_j^2 = -1$ shows that $\Theta_{i,j}$ has multiplicity 1 and cuts Θ_j at a rational point. Thus Θ_i has multiplicity 1 and is isomorphic to $\mathbb{P}_{k(s)}^1$. Otherwise, $\mathcal{Z}_0 \rightarrow \mathcal{Z}_i$ is an isomorphism in a neighborhood of $\Theta_{i,0}$. In particular, $\Theta_{i,0}$ is an exceptional divisor. Then we can conclude exactly as for Θ_0 . $\square$

Proposition 2.7. *Let $\mathcal{X}$ be a normal model of X over S . Then the following properties are equivalent:*

- (i) $\mathcal{X}$ is dominated by a semi-stable model over S ;
- (ii) X admits a semi-stable model over S and $\mathcal{X}_s$ is geometrically reduced for all $s \in S$;
- (iii) The minimal desingularization $\tilde{\mathcal{X}}$ of $\mathcal{X}$ is semi-stable over S .

Proof: (i) $\implies$ (ii) If $\mathcal{X}$ is dominated by a semi-stable model $\mathcal{X}'$, then any irreducible component Γ of $\mathcal{X}_s$ is birational to an irreducible component of $\mathcal{X}'_s$. The latter being geometrically reduced, $\mathcal{X}_s$ is geometrically reduced.

(ii) $\implies$ (iii) We know that $\tilde{\mathcal{X}}$ dominates a relatively minimal regular model $\mathcal{Z}$. The semi-stable reduction hypothesis implies that $\mathcal{Z}$ is semi-stable (see [13], 10.3.34(a) if $g(X) \geq 1$; if $g(X) \leq 0$, then $\mathcal{Z}$ is smooth). Since the minimal desingularization commutes with étale base change (see the proof of [13], Prop. 9.3.28), we can suppose that S is local with separably closed residue field. By Lemma 2.6, $\tilde{\mathcal{X}}_s$ is geometrically reduced. The map $\mathcal{X} \rightarrow \mathcal{Z}$ consists in blowing-up successively closed points. The fact that $\tilde{\mathcal{X}}_s$ is geometrically reduced implies that we only blow-up rational points in the smooth locus. Since $\mathcal{Z}$ is semi-stable, then so is $\mathcal{X}$. $\square$

Corollary 2.8. *There exists a finite separable extension K'/K such that $\mathcal{X}_{S'}$, where S' is the normalization of S in K' , is dominated by a semi-stable model of $X_{K'}$.*

Proof: We can suppose that X has semi-stable reduction over S . Since X has good reduction over an open dense subset of S , we can suppose that S is local. By the finiteness Theorem of Grauert-Remmert ([8], see also [3], Theorem 1.3) applied to the formal completion of $\mathcal{X}$ along its special fiber, there exists a finite Galois extension $L/\hat{K}$ such that the normalization of $\mathcal{X}_{\mathcal{O}_L}$ has geometrically reduced special fiber. See [7], p. 247 for how to descend the result to K (Note that [12] fills a gap in the proof of a main theorem in [7]). We then apply Proposition 2.7. $\square$

Remark 2.9 The corollary is useful in a recent work of Ch. Deninger and A. Werner on vector bundles and representations of the fundamental group of p -adic curves [6]. In fact, de Jong ([11], Theorem 2.4) already proved it in the situation when S is a Noetherian integral excellent scheme of any dimension, and when X is an integral curve over $K(S)$. The scheme S' is then proper and generically finite over S . The proof here for one-dimensional S is simpler, and more effective in some sense.

Remark 2.10 When S is local and complete, the corollary can be reformulated in terms of rigid analytic geometry as follows: let $\mathcal{U}$ be a formal covering of X . Then after finite separable extension of K , $\mathcal{U}$ can be refined to a distinguished formal covering $\mathcal{V}$ with semi-stable reduction. As such, the statement can be easily worked out using [1], Theorem 5.5, and step 2 in the proof of Lemma 7.3, page 377. The non complete case can then be obtained using Proposition 1.3.

Remark 2.11 (*Effective reduced fiber theorem*). In the case of 2.8, we can give an effective method to eliminate the multiplicities of $\mathcal{X}_s$, without using Grauert-Remmert's theorem. Suppose that X has semi-stable reduction. Let Γ be an irreducible component of $\mathcal{X}_s$ of geometric multiplicity $d > 1$. If we choose two closed points P_1, P_2 of X which specialize to two distinct points in the interior of Γ , and if we take $L = K(P_1, P_2)$, then the irreducible components of $(\mathcal{X}_{\mathcal{O}_L})'$'s (where $(\mathcal{X}_{\mathcal{O}_L})'$ denotes the normalization of $\mathcal{X}_{\mathcal{O}_L}$) lying above Γ are geometrically reduced. If K is strictly Henselian, we can bound $[L : K]$ by d^2 . Note that L can be chosen to be separable over K . If X has not necessarily semi-stable reduction, then we can bound $[L : K]$ by the max of d^2 and a constant depending only on g .

2.12 The stable hull. Now let us construct a minimal semi-stable model dominating $\mathcal{X}$. Let $\mathcal{Z}$ be a locally complete intersection (e.g., regular or semi-stable) model of X over S . Let $\omega_{\mathcal{Z}/S}$ be the (invertible) dualizing sheaf of $\mathcal{Z}/S$. Recall that a (-2) -curve on $\mathcal{Z}$ is an irreducible component Γ of a closed fiber $\mathcal{Z}_s$ such that $\deg \omega_{\mathcal{Z}/S}|_{\Gamma} = 0$. If $\mathcal{Z}$ is semi-stable and $k(s)$ is algebraically closed, and Γ is not a connected component of $\mathcal{Z}_s$, then this is equivalent to $\Gamma \simeq \mathbb{P}_{k(s)}^1$ and Γ meets the other irreducible components at exactly two points. Recall that the *exceptional locus* of a birational projective morphism $\pi : \mathcal{Z} \rightarrow \mathcal{X}$ is by definition the complementary of $\pi^{-1}(U)$, where U is the biggest open subscheme of $\mathcal{X}$ such that $\pi^{-1}(U) \rightarrow U$ is an isomorphism. When $\mathcal{X}$ is normal, the exceptional locus is equal to the union of the prime divisors of $\mathcal{Z}$ which map to closed points in $\mathcal{X}$. A semi-stable model $\mathcal{Z}$ dominating $\mathcal{X}$ will be called *relatively minimal* if there is no semi-stable model between $\mathcal{X}$ and $\mathcal{Z}$, except $\mathcal{Z}$ itself.

Lemma 2.13. *Let $\mathcal{Z}$ be a semi-stable model of X over S . Let V be an effective vertical divisor on $\mathcal{Z}$ such that for all $s \in S$, no connected component of $\mathcal{Z}_s$ is contained in V .*

- (a) *If $\deg \omega_{\mathcal{Z}/S}|_{\Gamma} \leq 0$ for all components Γ of V , then there exists a contraction map $\mathcal{Z} \rightarrow \mathcal{W}$ of V and $\mathcal{W}$ is semi-stable.*
- (b) *If there exists a contraction map $\mathcal{Z} \rightarrow \mathcal{W}$ of V with $\mathcal{W}$ semi-stable, then $\deg \omega_{\mathcal{Z}/S}|_{\Gamma} \leq 0$ for at least one irreducible component Γ of V .*

Proof: (a) is well-known but we were not able to find a proper reference. We can suppose that S is local. Let $\rho : \tilde{\mathcal{Z}} \rightarrow \mathcal{Z}$ be the minimal desingularization. Then the components Θ of the exceptional locus E of ρ are (-2) -curves. Let $\tilde{V}$ be the strict transform of V in $\tilde{\mathcal{Z}}$. Let us show that $V' := E + \tilde{V}$ can be contracted. We have $\omega_{\tilde{\mathcal{Z}}/S} = \rho^* \omega_{\mathcal{Z}/S}$ because $\mathcal{Z}$ is semi-stable. Hence $\deg \omega_{\tilde{\mathcal{Z}}/S}|_{\Gamma'} \leq 0$ for all components Γ' of V' . If there exists a Γ' such that $\deg \omega_{\tilde{\mathcal{Z}}/S}|_{\Gamma'} < 0$, then Γ' is an exceptional divisor. Let $\pi : \tilde{\mathcal{Z}} \rightarrow \mathcal{Z}'$ be the contraction of Γ' . Then $\omega_{\tilde{\mathcal{Z}}/S} = \pi^* \omega_{\mathcal{Z}'/S}(\Gamma')$. We deduce easily that $\deg \omega_{\mathcal{Z}'/S}|_{\Gamma''} \leq 0$ for all Γ'' in $\pi(V')$. So by successively blowing-down exceptional divisors, we can suppose that V' consists only of (-2) -curves. By Artin's criterion of contractibility ([13], 9.4.7), V' can be contracted. Therefore V can be contracted.

It remains to see that $\mathcal{W}$ is semi-stable. Let $\mathcal{O}_{K'}$ be a discrete valuation ring containing $\mathcal{O}_S$ and let $S' = \text{Spec } \mathcal{O}_{K'}$. It is enough to show that $\mathcal{W}_{S'}$ is semi-stable. The map $\mathcal{Z}_{S'} \rightarrow \mathcal{W}_{S'}$ is the contraction of $V_{k(s')}$. We have $\omega_{\mathcal{Z}_{S'}/S'} = \omega_{\mathcal{Z}/S} \otimes \mathcal{O}_{S'}$. If Γ' is a component of $\mathcal{Z}_{s'}$ lying over $\Gamma \subseteq V$, then

$$[k(s') : k(s)] \deg_{k(s')} \omega_{\mathcal{Z}_{S'}/S'}|_{\Gamma'} = [k(\Gamma') : k(\Gamma)] \deg_{k(s)} \omega_{\mathcal{Z}/S}|_{\Gamma} \leq 0.$$

So we can reduce the lemma to the case $k(s)$ algebraically closed. Let $\Gamma \subseteq V$. Then $p_a(\Gamma) = 0$, $\Gamma \simeq \mathbb{P}_{k(s)}^1$, and Γ meets the other components of $\mathcal{Z}_s$ in at most two points. Now it is well-known that $\mathcal{W}_s$ is semi-stable (see for instance [13], 10.3.31).

(b) The previous computations show that $\deg \omega_{\mathcal{Z}/S}|_{\Gamma} = \deg \omega_{\tilde{\mathcal{Z}}/S}|_{\tilde{\Gamma}}$ for any irreducible component Γ of $\mathcal{Z}_s$. Let $\tilde{\mathcal{W}}$ be the minimal desingularization of $\mathcal{W}$. Suppose that $\tilde{\mathcal{Z}} \rightarrow \tilde{\mathcal{W}}$ is not an isomorphism. Let Γ be a component of $\mathcal{Z}_s$ whose strict transform in $\tilde{\mathcal{Z}}$ is an exceptional divisor contracted into a closed point in $\tilde{\mathcal{W}}$. Then $\Gamma \subseteq V$ and $\deg \omega_{\mathcal{Z}/S}|_{\Gamma} = \deg \omega_{\tilde{\mathcal{Z}}/S}|_{\tilde{\Gamma}} < 0$. If $\tilde{\mathcal{Z}} = \tilde{\mathcal{W}}$, then $\deg \omega_{\mathcal{Z}/S}|_{\Gamma} = \deg \omega_{\tilde{\mathcal{W}}/S}|_{\tilde{\Gamma}} = 0$ for all Γ in V because $\tilde{\Gamma}$ (the strict transform of Γ) is a (-2) -curve in $\tilde{\mathcal{W}}$. $\square$

Proposition 2.14. *Let $\mathcal{X}$ be a model of X over S dominated by a semi-stable model.*

- (a) *A semi-stable model $\mathcal{Z}$ dominating $\mathcal{X}$ is relatively minimal if and only if for all irreducible components of the exceptional locus of $\mathcal{Z} \rightarrow \mathcal{X}$, we have $\deg \omega_{\mathcal{Z}/S}|_{\Gamma} > 0$.*
- (b) *Let $\tilde{\mathcal{X}} \rightarrow \mathcal{X}$ be the minimal desingularization of $\mathcal{X}$. Let $\tilde{\mathcal{X}} \rightarrow \mathcal{W}$ be the contraction of the (-2) -curves contained in the exceptional locus of $\tilde{\mathcal{X}} \rightarrow \mathcal{X}$. Then $\mathcal{W}$ is the stable hull of $\mathcal{X}$.*

Proof: (a) is an immediate consequence of Lemma 2.13.

(b) By 2.13(a), $\mathcal{W}$ is semi-stable and dominates $\mathcal{X}$. Let $\mathcal{Z}$ be a semi-stable model dominating $\mathcal{X}$, and relatively minimal. Let us first show that $\tilde{\mathcal{X}}$ dominates $\mathcal{Z}$. Let $\tilde{\mathcal{Z}}$ be the minimal desingularization of $\mathcal{Z}$. Then $\tilde{\mathcal{Z}}$ dominates $\tilde{\mathcal{X}}$. Suppose that $\tilde{\mathcal{Z}} \rightarrow \tilde{\mathcal{X}}$ is not an isomorphism. Let Θ be an exceptional divisor of $\tilde{\mathcal{Z}}$ mapped to a closed point of $\tilde{\mathcal{X}}$. Since $\tilde{\mathcal{Z}} \rightarrow \mathcal{Z}$ is minimal, Θ maps to an irreducible component Γ of $\mathcal{Z}_s$ which is then contained in the exceptional locus of $\mathcal{Z} \rightarrow \mathcal{X}$. We have $\deg \omega_{\mathcal{Z}/S}|_{\Gamma} = \deg \omega_{\tilde{\mathcal{Z}}/S}|_{\Theta} < 0$. Contradiction. Therefore $\tilde{\mathcal{Z}} \simeq \tilde{\mathcal{X}}$ and $\tilde{\mathcal{X}} \rightarrow \mathcal{Z}$ consists in contracting some (-2) -curves in $\mathcal{X}_s$. Hence $\mathcal{Z}$ dominates $\mathcal{W}$. $\square$

Remark 2.15 A *Du Val model* of X over S is a model $\mathcal{X}$ such that $\deg \omega_{\tilde{\mathcal{X}}/S}|_{\Gamma} = 0$ where $\tilde{\mathcal{X}} \rightarrow \mathcal{X}$ is the minimal desingularization and where Γ is any irreducible component of the exceptional locus of $\tilde{\mathcal{X}} \rightarrow \mathcal{X}$. The above results (2.13 and 2.14) still hold when “semi-stable” is replaced by “Du Val”, except the base change property. The point is that Du Val models do not commute with base change.

Remark 2.16 Let $\mathcal{W}$ be semi-stable and dominating $\mathcal{X}$. Then $\mathcal{W}$ is the stable hull of $\mathcal{X}$ if and only if $\text{Aut}_{\mathcal{X}_s}(\mathcal{W}_s)$ is finite for all $s \in S$.

2.17 Proof of Theorem 2.3. (a) is contained in Corollary 2.8 and Proposition 2.14. (b) It is enough to show that $\mathcal{W}_{S'}$ is relatively minimal. Let Γ' be an irreducible component of $\mathcal{W}_{S'}$ contained in the exceptional locus of $\mathcal{W}_{S'} \rightarrow \mathcal{X}_{S'}$. The image of Γ' in $\mathcal{W}$ is in the exceptional locus of $\mathcal{W} \rightarrow \mathcal{X}$. Similarly to the proof of Lemma 2.13(a), we have $\deg \omega_{\mathcal{W}_{S'}/S'}|_{\Gamma'} > 0$. Therefore $\mathcal{W}_{S'}$ is relatively minimal by 2.14(a). $\square$

Remark 2.18 Suppose that $g(X) \geq 1$ and S is affine. Let $\mathcal{X}$ be the minimal regular model of X over S and let $\mathcal{W}$ be the stable hull of $\mathcal{X}_{S'}$ over some extension S'/S . Let $\mathcal{X}'$ be the stable or minimal regular model of $X_{K'}$ over S' . Then $H^0(\mathcal{W}, \omega_{\mathcal{W}/S'}) = H^0(\mathcal{X}', \omega_{\mathcal{X}'/S'})$. One should be able to recover some arithmetic informations on $\mathcal{X}$ from the sheaf $\omega_{\mathcal{W}/S'} \otimes ((\omega_{\mathcal{X}/S})^\vee \otimes \mathcal{O}_{S'})$. Let us consider the

ideal

$$\mathcal{O}_{S'}(-\delta) := H^0(\mathcal{W}, \omega_{\mathcal{W}/S'}) \otimes (H^0(\mathcal{X}, \omega_{\mathcal{X}/S})^\vee \otimes \mathcal{O}_{S'}).$$

For example, if X is an elliptic curve over K , then for every closed point $s \in S$, we can show that

$$12\text{ord}_s(\delta) = \text{ord}_s(\Delta) + a_s \text{ord}_s(j),$$

where Δ is the minimal discriminant divisor of X over S , and $a_s = 0$ if X has potentially good reduction at s , $a_s = 1$ otherwise.

2.19 Marked curves. Recall that a (proper) *marked curve* $Z \rightarrow T$ over a scheme T is a proper flat scheme of relative dimension 1 over T endowed with a finite set $M \subset Z(T)$ of sections with pairwise disjoint supports contained in the smooth locus of Z/T (for our purpose, it is not necessary to order these sections). Note that if T is irreducible with generic point ξ , then M is determined by its generic fiber $M \cap Z_\xi$. We say that (Z, M) is *semi-stable* if $Z \rightarrow T$ is semi-stable. We say that (Z, M) is *stable* if it is semi-stable and if for any geometric point $\bar{t}$ of T , $Z_{\bar{t}}$ is connected and for any irreducible component Γ of $Z_{\bar{t}}$, Γ meets the other components in at least $1 - (2p_a(\Gamma) - 2) - |M \cap \Gamma|$ points. This amounts to say that $\omega_{Z/T}(M)$ is ample.

A *morphism* of marked curves $(Z, M) \rightarrow (Z', M')$ over T is a T -morphism $f : Z \rightarrow Z'$ such that $f(M) \subseteq M'$.

Let (X, M) be a smooth marked curve over $K = K(S)$. A *marked model* of (X, M) over S is a marked curve $(\mathcal{X}, \mathcal{M})$ over S whose generic fiber is isomorphic to (X, M) . Since $\mathcal{M}$ is uniquely determined by M and $\mathcal{X}$, we will omit $\mathcal{M}$ in the notation $(\mathcal{X}, \mathcal{M})$ and we will simply say $\mathcal{X}$ is a marked model of (X, M) .

Let $\mathcal{X}$ be a model of X over S . The *stable marked hull* of $\mathcal{X}$ is the minimal semi-stable marked model of (X, M) dominating $\mathcal{X}$. Note that a stable marked hull is not necessarily a stable marked curve.

Corollary 2.20. *Let (X, M) be a smooth marked curve over K and let $\mathcal{X}$ be a (non-marked) model of X . Then after finite separable extension of K , $\mathcal{X}$ admits a stable marked hull. More precisely, if $\mathcal{X}$ has a stable hull over some extension S'/S , then it has a stable marked hull over S' . Moreover, the formation of stable marked hull commutes with flat base change of Dedekind schemes.*

Proof: We can suppose that $\mathcal{X}$ has a stable hull $\mathcal{Z}$ over S . Let $\tilde{\mathcal{Z}}$ be a desingularization of $\mathcal{Z}$, let $\overline{M}$ be the Zariski closure of M in $\tilde{\mathcal{Z}}$ and let $\tilde{\mathcal{Z}}_{s_1}, \dots, \tilde{\mathcal{Z}}_{s_n}$ be the fibers such that $\overline{M} \rightarrow \overline{M} \cap \tilde{\mathcal{Z}}_{s_i}$ is not injective. Let $\mathcal{Z}' \rightarrow \tilde{\mathcal{Z}}$ be an embedded resolution $\overline{M} + \sum_i \tilde{\mathcal{Z}}_{s_i}$ in $\tilde{\mathcal{Z}}$ so that the Zariski closure $\mathcal{M}'$ of M in $\mathcal{Z}'$ is a disjoint union of sections (contained in the smooth locus because $\mathcal{Z}'$ is regular). Then $\mathcal{Z}'$ is a semi-stable marked model dominating $\mathcal{X}$.

Let $\mathcal{W}$ be any semi-stable marked model of (X, M) over S . Similarly to the non marked case, we can show that $\mathcal{W}$ is relatively minimal if and only if $\omega_{\mathcal{W}/S}(\mathcal{M})|_\Gamma$, where $\mathcal{M}$ is the Zariski closure of M in $\mathcal{W}$, has positive degree for all Γ in the exceptional locus of $\mathcal{W} \rightarrow \mathcal{X}$. This then implies that the stable marked hull is obtained by contracting prime divisors Γ in the exceptional locus of $\mathcal{Z}' \rightarrow \mathcal{Z}$ such that $\deg(\omega_{\mathcal{Z}'/\mathcal{Z}}(\mathcal{M}')|_\Gamma) \leq 0$, and that the stable marked hull commutes with flat base change. $\square$

Remark 2.21 If (X, M) is stable (meaning that $2g(X) - 2 + |M| \geq 1$) and if X has semi-stable reduction over S , then there exists a semi-stable marked model $\mathcal{X}$ of (X, M) over S and minimal for this property. This model is the stable marked model

of (X, M) . It is characterized by the property that for all irreducible components Γ of $\mathcal{X}_s$, one has $\deg \omega_{\mathcal{X}/S}(\mathcal{M})|_{\Gamma} > 0$, where $\mathcal{M}$ is the Zariski closure of M in $\mathcal{X}$. As above, this implies that the stable marked model commutes with base change.

3. SEMI-STABLE MODELS OF FINITE COVERS

We (re)prove that any finite morphism of projective smooth curves over K extends, after finite separable extension of K , to a finite morphism of semi-stable models.

Definition 3.1 Let $f : X \rightarrow Y$ be a finite morphism of smooth connected projective curves over $K(S)$. A *model (or extension) of f over S* consists in a morphism $\mathcal{X} \rightarrow \mathcal{Y}$ over S extending f , where $\mathcal{X}$ and $\mathcal{Y}$ are models over S of X and Y respectively. A model of f is said to be *finite* if it is a finite morphism, and *semi-stable* (see [4]) if it is finite and if $\mathcal{X}, \mathcal{Y}$ are semi-stable. We say that a model $\mathcal{X} \rightarrow \mathcal{Y}$ of f dominates another one $\mathcal{X}' \rightarrow \mathcal{Y}'$ if there are birational morphisms $\mathcal{X} \rightarrow \mathcal{X}'$, $\mathcal{Y} \rightarrow \mathcal{Y}'$ making the following diagram commutative

$$\begin{array}{ccc} \mathcal{X} & \longrightarrow & \mathcal{Y} \\ \downarrow & & \downarrow \\ \mathcal{X}' & \longrightarrow & \mathcal{Y}' \end{array}$$

A model of f is *stable* if it is semi-stable and minimal (for the domination relation) among semi-stable models of f . If $\mathcal{X}, \mathcal{Y}$ are respective models of X, Y . Then the semi-stable model $\mathcal{X}' \rightarrow \mathcal{Y}'$ of f such that $\mathcal{X}'$ dominates $\mathcal{X}$, $\mathcal{Y}'$ dominates $\mathcal{Y}$, and which is minimal for these property, is called the *stable hull* of (the rational map) $\mathcal{X} \dashrightarrow \mathcal{Y}$. We can obviously make similar definitions for marked curves.

Remark 3.2 For a given $f : X \rightarrow Y$, the semi-stable models are not unique: let $\mathcal{X} \rightarrow \mathcal{Y}$ be a semi-stable model of f , let $\mathcal{Y}' \rightarrow \mathcal{Y}$ be a blowing-up along a closed point, then the stable hull of $\mathcal{X} \dashrightarrow \mathcal{Y}'$ (see 4.5) is a new semi-stable model.

3.3 Decomposition of inseparable morphisms. Let us first deal with purely inseparable morphisms $X \rightarrow Y$. The next two statements are well-known at least over perfect base fields.

Lemma 3.4. *Let K be a field of characteristic $p > 0$. Let E/F be a finite extension of function fields of one variable over K , with E separable over K . Then there exists a unique purely inseparable sub-extension L/F of E/F such that E/L is separable. Moreover, $F = KL^{p^r}$ for some $r \geq 0$.*

Proof: Let F_s be the separable closure of F in E . By [13], Cor. 3.2.27 (here we use the hypothesis E separable over K), there exists $r \geq 0$ such that $F_s = KE^{p^r}$. Let

$$L := \{e \in E \mid e^{p^r} \in F\}.$$

Then L/F is a purely inseparable extension, $F = KL^{p^r}$, and E/L is separable because otherwise $L \subseteq KE^p$ and $F_s \subseteq KE^{p^{r+1}}$. The uniqueness of L is obvious because it is necessarily equal to the radical closure of F in E . $\square$

Proposition 3.5. *Let $f : X \rightarrow Y$ be a finite morphism of normal connected curves over a field K of characteristic $p > 0$. Suppose that X is smooth. Then f can be decomposed into a finite separable morphism $X \rightarrow Z$ followed by $Z \rightarrow Y$ which can be identified to a Frobenius map $Z \rightarrow Z^{(p^r)}$ for some $r \geq 0$. Moreover Z is smooth.*

Proof: Let L be the radicial closure of $K(Y)$ in $K(X)$, and let Z be the normalization of Y in L . Then f induces a finite separable morphism $X \rightarrow Z$. It is flat because Z is regular of dimension 1. Let $\bar{K}$ be an algebraic closure of K , then $X_{\bar{K}} \rightarrow Z_{\bar{K}}$ is flat, hence $Z_{\bar{K}}$ is regular. Finally, $Z \rightarrow Y$ can be identified to $Z \rightarrow Z^{(p^r)}$ by [13], 7.4.21. $\square$

Note that f can also be decomposed into $X \rightarrow X^{(p^r)}$ followed by a separable morphism $X^{(p^r)} \rightarrow Y$.

3.6 Semi-stable models.

Lemma 3.7. *Let $f_i : X \rightarrow Y_i$, $i = 1, \dots, n$ be finite surjective morphisms of integral projective varieties over K and let $\mathcal{Y}_i$ be a model of Y_i over S . Then there exists a model $\mathcal{X}$ of X over S such that $\mathcal{X}$ dominates $\mathcal{Y}_i$ (that is, $X \rightarrow Y_i$ extends to $\mathcal{X} \rightarrow \mathcal{Y}_i$) for all i .*

Proof: For N big enough, we have a closed immersion $g : X \rightarrow \mathbb{P}_K^N \times_K (\prod_{K,i} Y_i)$ induced by the projective morphism $\prod_i f_i : X \rightarrow \prod_{K,i} Y_i$. Now take $\mathcal{X}$ to be the Zariski closure of $g(X)$ in $\mathbb{P}_S^N \times_S (\prod_{S,i} \mathcal{Y}_i)$, endowed with the reduced structure. Note that if X is geometrically reduced, we can also use Lemma 4.3 with $\mathcal{X}_i = N(\mathcal{Y}_i, K(X))$. $\square$

Proposition 3.8. *Let S be a Dedekind scheme, and let $f : X \rightarrow Y$ be a finite morphism of smooth geometrically connected projective curves over $K := K(S)$. Let $\mathcal{X}, \mathcal{Y}$ be respective models of X and Y . Then there exists a finite separable extension K'/K such that over the normalization S' of S in K' , the cover $X_{K'} \rightarrow Y_{K'}$ extends to a finite morphism $\mathcal{X}' \rightarrow \mathcal{Y}'$, where $\mathcal{X}'$ (resp. $\mathcal{Y}'$) is a semi-stable model of X (resp. Y) over S dominating $\mathcal{X}_{S'}$ (resp. $\mathcal{Y}_{S'}$).*

Proof: Let $X \rightarrow Z \rightarrow Y$ be the decomposition given by Proposition 3.5. Let $\hat{X} \rightarrow Z$ be the Galois closure of $X \rightarrow Z$. After a finite separable extension of K , $\hat{X}$ is smooth over K . Let $\hat{\mathcal{X}}_0/S$ be a model of $\hat{X}$ dominating $\mathcal{X}$ and $\mathcal{Y}$ (3.7). Let $G := \text{Gal}(K(\hat{X})/K(Z))$. By 2.5, after a finite separable extension of K , there exists a semi-stable model $\hat{\mathcal{X}}$ of $\hat{X}$ endowed with an action of G and dominating $\hat{\mathcal{X}}_0$. Let $\mathcal{X}' = \hat{\mathcal{X}}/H$ where $H = \text{Gal}(K(\hat{X})/K(X))$, and $\mathcal{Z}' = \hat{\mathcal{X}}/G$. Then $\mathcal{X}' \rightarrow \mathcal{Z}'$ is a finite morphism of semi-stable models of X and Z respectively ([17], Prop. 5). Let $\mathcal{Y}' = \mathcal{Z}'^{(p^r)}$. Then the canonical map $\mathcal{Z}' \rightarrow \mathcal{Y}'$ is finite and $\mathcal{Y}'$ is semi-stable (loc. cit., or [13], Exerc. 10.3.19(a)). Since $\hat{\mathcal{X}}$ dominates $\mathcal{Y}$ and is finite over $\mathcal{Y}'$, we see easily that $\mathcal{Y}'$ dominates $\mathcal{Y}$ (use for instance [15], 4.1). Hence the proposition is proved with f' equal to the composition $\mathcal{X}' \rightarrow \mathcal{Z}' \rightarrow \mathcal{Y}'$. $\square$

Remark 3.9 If S is any Noetherian integral excellent scheme, then using the result of de Jong [11] as quoted in 2.9, we see that the proposition is still true.

The next corollary was known for separable morphisms ([4] when K is complete; [15], Remark 4.6 when $g(X) \geq 1$).

Corollary 3.10. *Let $f : X \rightarrow Y$ be a finite morphism of smooth geometrically connected projective curves over K . Then after a finite separable extension of K , there exists a finite morphism $\varphi : \mathcal{X} \rightarrow \mathcal{Y}$ of semi-stable models of X, Y respectively.*

Proof: Apply Proposition 3.8 to any pair of models of X, Y . $\square$

Lemma 3.11. *Let S be local. Let F be a finite closed subset of X . Let $\mathcal{Z}$ be a semi-stable model of X over S . Then there exists an integer $d > 0$ such that for any*

finite extension $\mathcal{O}_{K'}/\mathcal{O}_S$ of discrete valuation rings with ramification index divisible by d , if $\tilde{Z}'$ denotes a desingularization of $Z_{\mathcal{O}_{K'}}$, then the Zariski closure of $F_{K'}$ in $\tilde{Z}'$ is contained in the smooth locus of $\tilde{Z}'$.

Proof: Let $\alpha \in F$, and let $x \in \mathcal{Z}_s$ be a singular specialization of α . The local ring of an étale neighborhood of $x \in \mathcal{Z}$ is isomorphic to $\mathcal{O}_K[[u, v]]/(uv - a)$, with a a power of a uniformizing element of $\mathcal{O}_K$. Let $u(\alpha)$ be the image of u in $K(\alpha)$. Then $|a| < |u(\alpha)| < 1$. After an extension of big enough ramification index, $|u(\alpha)|$ belongs to $|K|$. If this condition is satisfied for all $\alpha \in F$ and for all singular specializations of α , then it is easy to see that the specializations of F in $\tilde{Z}'$ are smooth points. Indeed, if $\mathcal{Z}$ is regular, the parameter a in the above local ring is a uniformizing element, hence $|a| < |u(\alpha)| < 1$ cannot hold in $|K|$, so F must specialize to smooth points. $\square$

Proposition 3.12. *Let $f : (X, M) \rightarrow (Y, N)$ be a finite morphism of smooth geometrically connected marked projective curves over K , let $\mathcal{X}, \mathcal{Y}$ be respective models of X, Y . Then after a finite separable extension of K , there exists a semi-stable marked model $\mathcal{X}'$ (resp. $\mathcal{Y}'$) of (X, M) (resp. (Y, N)) such that $\mathcal{X}'$ and $\mathcal{Y}'$ dominate respectively $\mathcal{X}$ and $\mathcal{Y}$, and f extends to a finite morphism $\mathcal{X}' \rightarrow \mathcal{Y}'$.*

Proof: After enlarging K and replacing $\mathcal{X}$ and $\mathcal{Y}$ by their respective stable marked hull, we can suppose that $\mathcal{X}, \mathcal{Y}$ are semi-stable and that the Zariski closure $\overline{M}$ of M in $\mathcal{X}$ is a disjoint union of sections, and the same for N in $\mathcal{Y}$. Let $X \rightarrow Z \rightarrow Y$ be the decomposition as given by Proposition 3.5, and let $\hat{X} \rightarrow Z$ be the Galois closure of $X \rightarrow Z$. Let $f : \hat{X} \rightarrow X$ and $g : \hat{X} \rightarrow Y$ be the canonical morphisms. By lemma 3.11, after a finite separable extension, and after replacing $\hat{X}$ by its minimal desingularization (the group G still acts), we can suppose that the Zariski closure of $f^{-1}(M) \cup g^{-1}(N)$ in $\hat{X}$ is contained in the smooth locus. Then the Zariski closure of M in $\mathcal{X}'$ is contained in the smooth locus because $\hat{\mathcal{X}}_{\text{sm}}/\overline{H}$ is smooth, and it is a disjoint union of sections because $\mathcal{X}'$ dominates $\mathcal{X}$ and $\overline{M}$ is already a disjoint union of sections. Hence $\mathcal{X}'$ is semi-stable marked for (X, M) . The same arguments hold for $\mathcal{Y}'$. $\square$

4. STABLE HULL OF A MORPHISM

Definition 4.1 Let $f : X \rightarrow Y$ be a finite morphism of connected smooth projective curves over K . Let $\mathcal{X}, \mathcal{Y}$ be respective models of X, Y over S . The *finite hull* of $\mathcal{X} \dashrightarrow \mathcal{Y}$ is a finite model $\mathcal{X}^f \rightarrow \mathcal{Y}^f$ of f over S , such that $\mathcal{X}^f$ and $\mathcal{Y}^f$ are normal models of X, Y dominating respectively $\mathcal{X}$ and $\mathcal{Y}$, and which is minimal (for the domination relation) with respect to these properties.

Lemma 4.2. *Let $\mathcal{Y}$ be an integral scheme locally of finite type over S , let L be a finite extension of $K(\mathcal{Y})$, separable over $K = K(S)$, and let $\mathcal{X}$ be the normalization of $\mathcal{Y}$ in L . Then $\mathcal{X}$ is finite over $\mathcal{Y}$.*

Proof: The assertion is of course trivial if S is excellent. Let $y \in \mathcal{Y}_s$, let $A = \mathcal{O}_{\mathcal{Y}, y}$, and let B be the integral closure of A in L . We have to show that B is finite over A . Let $\mathcal{O}_K = \mathcal{O}_{S, s}$. Let C be the integral closure of $A \otimes_{\mathcal{O}_K} \hat{\mathcal{O}}_K$ in $L \otimes_K \hat{K}$. The latter is reduced (because L is separable over K) and finite over $K(\mathcal{Y}) \otimes_K \hat{K}$, the total ring of fractions of $A \otimes \hat{\mathcal{O}}_K$. Since $\hat{\mathcal{O}}_K$ is excellent, C (and thus $B \otimes \hat{\mathcal{O}}_K$) is finitely generated over $A \otimes \hat{\mathcal{O}}_K$. Then Nakayama's lemma implies that $B \otimes \hat{\mathcal{O}}_K$ is

generated over $A \otimes \hat{\mathcal{O}}_K$ by finitely many elements of B . These elements generate B over A because $\mathcal{O}_K \rightarrow \hat{\mathcal{O}}_K$ is faithfully flat.

Note that the proof still work if S is any Noetherian integral scheme such that $\hat{\mathcal{O}}_{S,s}$ is reduced for all $s \in S$. $\square$

Lemma 4.3. *Let X be an integral projective variety over K , and let $\mathcal{X}_1, \mathcal{X}_2, \dots, \mathcal{X}_n$ be models of X over S .*

- (a) *There exists a smallest model $\mathcal{X}$ of X dominating $\mathcal{X}_i$ for all i . Let us denote $\mathcal{X}$ by $\mathcal{X}_1 \vee \dots \vee \mathcal{X}_n$.*
- (b) *If X is geometrically integral, then for any flat morphism of Dedekind schemes $S' \rightarrow S$, we have $(\mathcal{X}_1 \vee \dots \vee \mathcal{X}_n)_{S'} = (\mathcal{X}_1)_{S'} \vee \dots \vee (\mathcal{X}_n)_{S'}$.*
- (c) *If $\dim X = 1$, then every irreducible component of $(\mathcal{X}_1 \vee \dots \vee \mathcal{X}_n)_s$ dominates an irreducible component of $(\mathcal{X}_i)_s$ for some i .*

Proof: The proof is similar to that of 2.4. Let $\mathcal{P}$ be the fiber product $\prod_{S,i} \mathcal{X}_i$ over S . Then the diagonal map makes X a closed subscheme of $\mathcal{P}_K$. Let $\mathcal{X}$ be the Zariski closure of X in $\mathcal{P}$ endowed with the reduced structure. Then $\mathcal{X}$ is a model of X over S dominating the $\mathcal{X}_i$'s. Let $\mathcal{Z}$ be a model of X over S dominating the $\mathcal{X}_i$'s. Then we have a natural morphism $f : \mathcal{Z} \rightarrow \mathcal{P}$ whose image $f(\mathcal{Z})$ is irreducible, with generic fiber $\mathcal{X}_K$. Hence $f(\mathcal{Z}) = \mathcal{X}$ and f factorizes through $\mathcal{Z} \rightarrow \mathcal{X} \rightarrow \mathcal{P}$. So $\mathcal{X}$ is minimal. If X is geometrically integral, then $\mathcal{X}_{S'}$ is an integral closed subscheme of $\mathcal{P}' := \prod_{S',i} (\mathcal{X}_i)_{S'}$, with generic fiber isomorphic to the diagonal of $\mathcal{P}'_K$. By construction, $\mathcal{X}_{S'}$ is equal to $(\mathcal{X}_1)_{S'} \vee \dots \vee (\mathcal{X}_n)_{S'}$.

Let $s \in S$. Then $\mathcal{X}_s$ is a closed subscheme of $\mathcal{P}_s = \prod_{k(s),i} (\mathcal{X}_i)_s$, pure of dimension $\dim X$. Let Γ be an irreducible component of $\mathcal{X}_s$. If $\dim X > 0$, then the image of Γ in $\mathcal{X}_{i_0}$ has positive dimension for some i_0 . If $\dim X = 1$, then the image of Γ in $(\mathcal{X}_{i_0})_s$ is an irreducible component. $\square$

Proposition 4.4. *Let $\mathcal{X} \dashrightarrow \mathcal{Y}$ be a rational map. Suppose that either S is local and Henselian, or $\mathcal{X}$ and $\mathcal{Y}$ are semi-stable or regular. Then the finite hull of $\mathcal{X} \dashrightarrow \mathcal{Y}$ exists. It commutes with base changes in the following sense : let $S' \rightarrow S$ be a flat morphism of Dedekind schemes. Then the finite hull of $\mathcal{X}_{S'} \dashrightarrow \mathcal{Y}_{S'}$ exists and is equal to $((\mathcal{X}^f)_{S'})^\sim \rightarrow ((\mathcal{Y}^f)_{S'})^\sim$, where $\sim$ means normalization.*

Proof: The rational map $\mathcal{X} \dashrightarrow \mathcal{Y}$ is defined and finite above an open dense subset of S . So we can suppose that $S = \text{Spec } \mathcal{O}_K$ is local with closed point s . The case when $\mathcal{X}, \mathcal{Y}$ are semi-stable or regular is easily reduced to the Henselian case by passing to the completion of $\mathcal{O}_K$ and using Proposition 1.3. Suppose that $\mathcal{O}_K$ is Henselian. Let $\mathcal{X}_1$ be the normalization of $\mathcal{X} \vee N(\mathcal{Y}, K(X))$ (see 4.3). We have a morphism $\mathcal{X}_1 \rightarrow \mathcal{Y}$. By [15], Lemma 4.14, there exists a (unique) normal model $\mathcal{Y}^f$ such that the rational map $\mathcal{X}_1 \dashrightarrow \mathcal{Y}^f$ is quasi-finite and surjective in codimension 1 (in other words : if $\mathcal{U}$ is the domain of definition of $\mathcal{X}_1 \dashrightarrow \mathcal{Y}^f$, then $\mathcal{U}_s$ is dense in $(\mathcal{X}_1)_s$, $\mathcal{U}_s \rightarrow \mathcal{Y}_s$ is quasi-finite and has dense image). Since $\mathcal{X}_1 \rightarrow \mathcal{Y}$ is a morphism, $\mathcal{Y}^f$ dominates $\mathcal{Y}$. Let $\mathcal{X}^f$ be the normalization of $\mathcal{Y}^f$ in $K(X)$. Then it is easy to see that $\mathcal{X}^f \rightarrow \mathcal{Y}^f$ is the finite hull of $\mathcal{X} \dashrightarrow \mathcal{Y}$ (use [15], 4.1 for instance).

It remains to prove the base change property. The rational map $(\mathcal{X}_{S'})^\sim \dashrightarrow ((\mathcal{Y}^f)_{S'})^\sim$ is quasi-finite and surjective in codimension 1. By the above construction, $(\mathcal{Y}_{S'})^f = ((\mathcal{Y}^f)_{S'})^\sim$ and $(\mathcal{X}_{S'})^f$ is the normalization of $((\mathcal{Y}^f)_{S'})^\sim$ in $K(X_{K(S')})$. Since $((\mathcal{X}^f)_{S'})^\sim \rightarrow ((\mathcal{Y}^f)_{S'})^\sim$ is finite, $(\mathcal{X}_{S'})^f$ is isomorphic to $((\mathcal{X}^f)_{S'})^\sim$. $\square$

Theorem 4.5. *Let S be a connected Noetherian regular scheme of dimension 1, let $f : X \rightarrow Y$ be a finite morphism of smooth geometrically connected projective curves over $K := K(S)$. Let $\mathcal{X}, \mathcal{Y}$ be respective models of X, Y over S . Then after a finite separable extension of K , $\mathcal{X} \dashrightarrow \mathcal{Y}$ admits a stable hull $\mathcal{X}' \rightarrow \mathcal{Y}'$. Moreover, the formation of $\mathcal{X}' \rightarrow \mathcal{Y}'$ commutes with flat base change.*


Proof: After a finite separable extension of K , we can suppose that there exists a semi-stable model $\mathcal{X}_\infty \rightarrow \mathcal{Y}_\infty$ of f dominating $\mathcal{X} \dashrightarrow \mathcal{Y}$ (Cor. 3.10). Let us show that $\mathcal{X} \dashrightarrow \mathcal{Y}$ then admits a stable hull over S . Consider the stable hull $\mathcal{Y}_1$ of $\mathcal{Y}$ and the stable hull $\mathcal{X}_1$ of $N(\mathcal{Y}_1, K(X))$. Then $\mathcal{X}_1$ and $\mathcal{Y}_1$ are dominated by $\mathcal{X}_\infty$ and $\mathcal{Y}_\infty$ respectively. Let $\mathcal{X}_2 \rightarrow \mathcal{Y}_2$ be the finite hull of $\mathcal{X}_1 \rightarrow \mathcal{Y}_1$. Then it is also dominated by $\mathcal{X}_\infty \rightarrow \mathcal{Y}_\infty$. Now restart again the process of taking stable hull and finite hull with $\mathcal{X}_2 \rightarrow \mathcal{Y}_2$. We construct in this way an increasing sequence of (normal) models $\mathcal{X}_n \rightarrow \mathcal{Y}_n$ of f over S which are dominated by $\mathcal{X}_\infty \rightarrow \mathcal{Y}_\infty$. This sequence is stationary at some rank n_0 . Then $\mathcal{X}_{n_0} \rightarrow \mathcal{Y}_{n_0}$ is a semi-stable model of f . Note that the construction of $\mathcal{X}_n \rightarrow \mathcal{Y}_n$ does not depend on the choice of $\mathcal{X}_\infty \rightarrow \mathcal{Y}_\infty$. In particular, $\mathcal{X}_n \rightarrow \mathcal{Y}_n$ is dominated by any semi-stable model of f dominating $\mathcal{X} \dashrightarrow \mathcal{Y}$. Therefore, $\mathcal{X}_{n_0} \rightarrow \mathcal{Y}_{n_0}$ is the stable hull of $\mathcal{X} \dashrightarrow \mathcal{Y}$. Finally, the formation of the stable hull commutes with flat base change because the stable hull of a model and the finite hull of a morphism commute with flat base change (2.3, 4.4). $\square$

Corollary 4.6. *Suppose that either $g(X) \geq 2$, or $g(X) = 1$ and X has potentially good reduction. Then there exists a finite separable extension of K' of K such that $X_{K'} \rightarrow Y_{K'}$ admits a stable model $\mathcal{X}' \rightarrow \mathcal{Y}'$ over S' , where S' is the integral closure of S in K' . Moreover, for any Dedekind scheme T dominating S' , $\mathcal{X}'_T \rightarrow \mathcal{Y}'_T$ is the stable model of $X_{K(T)} \rightarrow Y_{K(T)}$.*

Proof: We can suppose that X has semi-stable reduction over S . The cover $X \rightarrow Y$ extends to a finite morphism of smooth projective models over a dense open subset of S . So we can suppose that $S = \text{Spec } \mathcal{O}_K$ is local. Let $\mathcal{X}^{\text{st}}$ be the stable (resp. smooth projective) model of X if $g(X) \geq 2$ (resp. if $g(X) = 1$). Suppose first that $\mathcal{O}_K$ is complete (hence Henselian). Let $\mathcal{X}^{\text{st}} \dashrightarrow \mathcal{Y}''$ be the rational map extending $X \rightarrow Y$ and which is quasi-finite and surjective in codimension 1 (see [15], 4.14). Then the stable hull $\mathcal{X} \rightarrow \mathcal{Y}$ of $\mathcal{X}^{\text{st}} \dashrightarrow \mathcal{Y}''$ is clearly the stable model of $X \rightarrow Y$. The construction of $\mathcal{X} \rightarrow \mathcal{Y}$ commutes with flat base change because that of $\mathcal{Y}''$ and the stable hull commute with flat base change. If $\mathcal{O}_K$ is non necessarily complete, we can use Corollary 1.6. $\square$

Remark 4.7 Let $X \rightarrow Y$ be as above. If $X \rightarrow Y$ has a semi-stable model over S , then it has a stable model over S . This can be seen in the proof of 4.6. If $X \rightarrow Y$ is moreover Galois of group G , and if $\mathcal{X}$ is the stable model (or smooth model if $g(X) = 1$) of X over S . Then the stable model of $X \rightarrow Y$ is equal to $\mathcal{X} \rightarrow \mathcal{X}/G$.

Remark 4.8 Suppose moreover that $X \rightarrow Y$ is separable, and that the Galois closure $\hat{X}$ of $X \rightarrow Y$ is smooth and geometrically connected over K (which is true after a finite separable extension of K). Let $\hat{\mathcal{X}}$ be the stable model of $\hat{X}$, and let $G = \text{Gal}(K(\hat{X})/K(Y))$, $H = \text{Gal}(K(\hat{X})/K(X))$ as in the proof of 3.8. Then we can ask whether $\hat{\mathcal{X}}/H \rightarrow \hat{\mathcal{X}}/G$ is the stable model of $X \rightarrow Y$. The answer is no in general. Let us give an example with X and Y having good reduction.



Let S be local, complete, with algebraically closed residue field k . Let $C_1 \rightarrow D_1$ be a finite separable morphism of degree $d \geq 3$ with $C_1, D_1 \simeq \mathbb{P}_k^1$, totally ramified above some point $y_1 \in D_1$, and such that the Galois closure E of $C_1 \rightarrow D_1$ is a curve of genus $g(E) \geq 1$. Let $C_2 \rightarrow D_2$ be a finite separable morphism of degree d of smooth connected projective curves over k , totally ramified above a point $y_2 \in D_2$ and such that $g(C_2) \geq 1$. Let D be the semi-stable curve over k obtained by identifying y_1 and y_2 . Let C be the semi-stable curve defined in a similar way. Then we have a finite morphism $\rho : C \rightarrow D$ which is generically étale, and such that $C_1 \rightarrow D_1, C_2 \rightarrow D_2$ have the same ramification index at y_1 and y_2 . By [14], Prop. 5.4, the cover $C \rightarrow D$ lifts to a finite morphism $\mathcal{C} \rightarrow \mathcal{D}$ over S with smooth generic fibers X, Y . Let $\mathcal{C} \rightarrow \mathcal{C}_2$ (resp. $\mathcal{D} \rightarrow \mathcal{D}_2$) be the contraction of C_1 (resp. of D_1). Then $\mathcal{C}_2, \mathcal{D}_2$ are smooth, and the canonical morphism $\mathcal{C}_2 \rightarrow \mathcal{D}_2$ is the stable model of $X \rightarrow Y$. Let $\mathcal{Z}$ be the normalization of $\mathcal{D}$ in $K(\hat{X})$. Let Θ be an irreducible component of $\mathcal{Z}_s$ lying over D_1 . Then the separable closure of $k(D_1)$ in $k(\Theta)$ is Galois over $k(D_1)$ ([18], I, §7, Prop. 20) and contains $k(C_1)$. Thus $k(\Theta)$ contains a subfield isomorphic to $k(E)$. In particular, $p_a(\Theta) \geq 1$. If $\mathcal{C}_2 \rightarrow \mathcal{D}_2$ is equal to $\hat{\mathcal{X}}/H \rightarrow \hat{\mathcal{X}}/G$, then $\mathcal{Z}$ dominates $\hat{\mathcal{X}}$ because $\mathcal{D}$ dominates $\mathcal{D}_2$. But Θ maps to a closed point of $\hat{\mathcal{X}}_s$, thus $\hat{\mathcal{X}}$ can not be semi-stable. Contradiction.

Remark 4.9 If X has genus 1 and multiplicative reduction at some point of S , or if $g(X) = 0$, then over any ramified extension of S , there is no stable model of the identity map $X \rightarrow X$. The reason we take a ramified extension S'/S is, when $g(X) = 1$, $X_{K'}$ has no minimal semi-stable model (see the proof of Lemma 1.5).

Let us give a characterization of the stable model.

Definition 4.10 Let $X_1 \rightarrow Y_1, X_2 \rightarrow Y_2$ be morphisms of schemes over some base scheme T . An *isomorphism* of the pairs $(X_1 \rightarrow Y_1) \rightarrow (X_2 \rightarrow Y_2)$ is given by T -isomorphisms $X_1 \rightarrow X_2, Y_1 \rightarrow Y_2$ such that the diagram

$$\begin{array}{ccc}
 X_1 & \longrightarrow & X_2 \\
 \downarrow & & \downarrow \\
 Y_1 & \longrightarrow & Y_2
 \end{array}$$

is commutative. We denote by $\text{Isom}_T((X_1 \rightarrow Y_1), (X_2 \rightarrow Y_2))$ the set of these isomorphisms. Now $\text{Aut}_T(X \rightarrow Y)$ has an obvious meaning.

Proposition 4.11. *Keep the hypothesis of 4.5 and suppose that $g(X) \geq 2$. Let $\mathcal{X} \rightarrow \mathcal{Y}$ be a semi-stable model of $X \rightarrow Y$. Consider the following properties.*

- (i) $\mathcal{X} \rightarrow \mathcal{Y}$ is the stable model of $X \rightarrow Y$ over S ;
- (ii) Let Γ be any irreducible component of $\mathcal{Y}_s$ such that $\deg \omega_{\mathcal{Y}_s/k(s)}|_{\Gamma} \leq 0$, then there exists an irreducible component Θ of $\mathcal{X}_s$ dominating Γ such that $\deg \omega_{\mathcal{X}_s/k(s)}|_{\Theta} > 0$;
- (iii) $\text{Aut}_{k(\bar{s})}(\mathcal{X}_{\bar{s}} \rightarrow \mathcal{Y}_{\bar{s}})$ is finite for all $s \in S$.

Then (i) $\iff$ (ii) $\implies$ (iii).

Proof: Looking in the proofs of 4.5 and 4.6, we see that $X \rightarrow Y$ admits a stable model over S . Thus $\mathcal{X} \rightarrow \mathcal{Y}$ is stable if and only if it is a relatively minimal semi-stable model. Hence the equivalence (i) $\iff$ (ii) is an immediate consequence of Lemma 2.13.

Suppose that Condition (ii) is satisfied. Then the same condition holds over $k(\bar{s})$ (with dualizing sheaves on $\mathcal{X}_{\bar{s}}$ and $\mathcal{Y}_{\bar{s}}$). Consider the natural inclusion

$$\mathrm{Aut}_{k(\bar{s})}(\mathcal{X}_{\bar{s}} \rightarrow \mathcal{Y}_{\bar{s}}) \subseteq \mathrm{Aut}_{k(\bar{s})}(\mathcal{X}_{\bar{s}}) \times \mathrm{Aut}_{k(\bar{s})}(\mathcal{Y}_{\bar{s}}).$$

Note that the right-hand side is not a finite group in general. Let G be the subgroup (of finite index) of $\mathrm{Aut}_{k(\bar{s})}(\mathcal{X}_{\bar{s}})$ consisting in automorphisms which fix globally each irreducible component, and let H be the subgroup of $\mathrm{Aut}_{k(\bar{s})}(\mathcal{Y}_{\bar{s}})$ consisting in automorphisms τ such that $\tau|_{\Gamma} = \mathrm{Id}$ for every irreducible component Γ with $\deg \omega_{\mathcal{Y}_{k(\bar{s})/k(\bar{s})}}|_{\Gamma} > 0$. Then H is of finite index in $\mathrm{Aut}_{k(\bar{s})}(\mathcal{Y}_{\bar{s}})$ because $\{\tau \in \mathrm{Aut}_{k(\bar{s})}(\mathcal{Y}_{k(\bar{s})}) \mid \tau(\Gamma) = \Gamma\}$ is finite for any such Γ . Thus it is enough to show that $G' := \mathrm{Aut}_{k(\bar{s})}(\mathcal{X}_{\bar{s}} \rightarrow \mathcal{Y}_{\bar{s}}) \cap (G \times H)$ is finite. Let I be the set of irreducible components Θ of $\mathcal{X}_{\bar{s}}$ such that $\deg \omega_{\mathcal{X}_{\bar{s}}/k(\bar{s})}|_{\Theta} > 0$. Then $G|_{\Theta}$ is finite for all $\Theta \in I$. Let (σ, τ) be an element of $\mathrm{Ker}(G' \rightarrow G \times H \rightarrow \prod_{\Theta \in I} G|_{\Theta})$. Condition (ii) implies that $\tau = \mathrm{Id}$ on $\mathcal{Y}_{\bar{s}}$, hence $\sigma \in \mathrm{Aut}_{\mathcal{Y}_{\bar{s}}}(\mathcal{X}_{\bar{s}})$. The latter is a finite group because $\mathcal{X}_{\bar{s}} \rightarrow \mathcal{Y}_{\bar{s}}$ is a finite morphism. Since the projection map $\mathrm{Aut}_{k(\bar{s})}(\mathcal{X}_{\bar{s}} \rightarrow \mathcal{Y}_{\bar{s}}) \rightarrow \mathrm{Aut}_{k(\bar{s})}(\mathcal{X}_{\bar{s}})$ is injective, the above kernel is finite. Hence G' is finite. $\square$

Remark 4.12 In general, (iii) does not imply (i). Let S be local, complete with algebraically closed residue field k . Let $\pi : \mathbb{A}_k^1 \rightarrow \mathbb{A}_k^1$ be a finite separable cover with trivial automorphisms group $\mathrm{Aut}_k(\pi)$. Then π extends to a finite cover $\mathbb{P}_k^1 \rightarrow \mathbb{P}_k^1$ totally ramified at ∞ . We can glue π with a finite separable cover $C_2 \rightarrow D_2$ of smooth projective curves of genus ≥ 2 over k and obtain a finite cover $C \rightarrow D$ with finite automorphisms group (see the construction in 4.8) which lifts to a finite morphism of semi-stable curves $\mathcal{C} \rightarrow \mathcal{D}$ over S . By the equivalence of (i) and (ii), $\mathcal{C} \rightarrow \mathcal{D}$ is not stable.

Remark 4.13 We can see that Theorem 4.5 holds for finite morphisms of smooth projective marked curves $(X, M) \rightarrow (Y, N)$: let $\mathcal{X}, \mathcal{Y}$ be respective models of X, Y , then after finite separable extension, there exists a stable marked hull $\mathcal{X}' \rightarrow \mathcal{Y}'$ of $\mathcal{X} \dashrightarrow \mathcal{Y}$. Moreover, the formation of $\mathcal{X}' \rightarrow \mathcal{Y}'$ commutes with flat base change. The proof is the same as for 4.5, except that we replace stable hull of a model by its stable marked hull, and we use 3.12 instead of 3.10.

The next lemma is a generalization of [15], 4.4(a) (take $N = \emptyset$).

Lemma 4.14. *Let $f : (X, M) \rightarrow (Y, N)$ be a finite morphism of connected smooth projective marked curves over K . Suppose that $2g(Y) - 2 + |N| > 1$ and $f^{-1}(N) = M$. If (X, M) has a stable marked model $\mathcal{X}$ over S , then (Y, N) has a stable marked model $\mathcal{Y}$, and $\mathcal{X} \dashrightarrow \mathcal{Y}$ is a morphism.*

Proof: The stable marked model exists over S when X has semi-stable reduction over S . So the existence of $\mathcal{X}$ implies that of $\mathcal{Y}$ (see [15], 4.8 when $g(Y) \geq 1$, the case $g(Y) = 0$ is trivial). It remains to show that the rational map $\mathcal{X} \dashrightarrow \mathcal{Y}$ is defined everywhere. Since the stable marked model commutes with flat base change (2.21), we can suppose that S is local with algebraically closed residue field and that $\mathcal{W} := N(\mathcal{Y}, K(X))$ has a stable marked hull $\mathcal{Z}$ over S . By definition, $\mathcal{Z}$ dominates $\mathcal{X}$. We are going to show that $\mathcal{Z} \rightarrow \mathcal{X}$ is an isomorphism. Or equivalently, that $\mathcal{Z}$ is the stable marked hull of $\mathcal{X}$. Let $\bar{M}$ denote the Zariski closure of M in $\mathcal{Z}$. It

is enough to show that $\deg \omega_{\mathcal{Z}/S}(\overline{M})|_{\Theta} > 0$ for all irreducible component Θ of $\mathcal{Z}_s$. If Θ is in the exceptional locus of $\mathcal{Z} \rightarrow \mathcal{W}$, then this is true because $\mathcal{Z} \rightarrow \mathcal{W}$ is the stable marked hull. Suppose that Θ is the strict transform of some irreducible component Γ of $\mathcal{W}_s$. Let Δ be the image of Γ in $\mathcal{Y}_s$. Then every point $y \in \Delta$ which is either a singular point of $\mathcal{Y}_s$ or a specialization of N lifts to a point of Θ which is either a singular point or a specialization of $f^{-1}(N) = M$ ([15], 4.4(b) for singular points; use going-down property as in [15], 4.3 for specializations of N). Since $\mathcal{Y}$ is stable marked, this implies that Θ contains at least three points of $(\mathcal{Z}_s)_{\text{sing}} \cup \overline{M}_s$, hence $\deg \omega_{\mathcal{Z}/S}(\overline{M})|_{\Theta} > 0$. $\square$

Remark 4.15 It is known that in general, $f : X \rightarrow Y$ does not extend to a morphism of the respective minimal regular models of X and Y over S (see e.g. [5], p. 333). However, if $g(Y) > 0$, then f extends to a morphism of the respective minimal regular models with normal crossings, at least if $k(s)$ is perfect for all closed points $s \in S$.

Corollary 4.16. *Let $f : (X, M) \rightarrow (Y, N)$ be a finite morphism of geometrically connected smooth projective marked curves over K . Suppose that $2g(Y) - 2 + |N| > 1$ and $f^{-1}(N) = M$. Then, after a finite separable extension of K , f admits a stable marked model $\mathcal{X} \rightarrow \mathcal{Y}$ over S . This construction commutes with flat base change.*

Remark 4.17 Let $f : X \rightarrow Y$ be a finite *separable* morphism of smooth projective curves. A natural way to mark X and Y is to take M equal to the ramification locus of f and N equal to the branch locus. By definition $f^{-1}(N) = M$. Of course, in general M is not contained in $X(K)$. But if f is tamely ramified (e.g., if $\text{char}(K) = 0$), then this becomes true over a finite separable extension of K ([15], 3.3). Moreover, if $g(Y) \geq 2$, or $g(Y) = 1$ and f is not étale, or if $g(Y) = 0$ and $g(X) \geq 1$, then $2g(Y) - 2 + |N| > 1$. So after again a finite separable extension of K , we have a canonical way to define a minimal semi-stable reduction of $X \rightarrow Y$ in which the (horizontal) ramification and branch loci are finite unions of sections contained in the smooth locus. If $(\deg f)!$ is invertible in $\mathcal{O}_S$, then $\mathcal{X}$ and $\mathcal{Y}$ are the respective stable marked models of X and Y ([16], §3.11, second Lemma).

Remark 4.18 Let $\hat{X}$, G , H be as in Remark 4.8. Let $\hat{X}$ be marked with its ramification locus over X (rational over K after a finite extension of K). Let $\hat{\mathcal{X}}$ be its stable marked model. If G has order prime to $\text{char}(k)$, then the stable marked model of $X \rightarrow Y$ is equal to $\hat{\mathcal{X}}/H \rightarrow \hat{\mathcal{X}}/G$ because both sides are respectively stable marked models of X and Y (see 4.17). But this is false if $\text{char}(k)$ divides $|G|$. Let us go back to the example of 4.8 and let $C_1 \rightarrow D_1$ be moreover étale outside of y_1 . Let $\mathcal{C}' \rightarrow \mathcal{C}$, $\mathcal{D}' \rightarrow \mathcal{D}$ be the stable marked hulls of $\mathcal{C}$, $\mathcal{D}$ (marked with horizontal ramification/branch loci). Let $\mathcal{C}' \rightarrow \mathcal{C}''$, $\mathcal{D}' \rightarrow \mathcal{D}''$ be the contraction of (the strict transforms of) C_1 , D_1 respectively. Then it is easy to see that the stable marked model (4.16) of $X \rightarrow Y$ is $\mathcal{C}'' \rightarrow \mathcal{D}''$. Similarly to 4.8, we see that it is different from $\hat{\mathcal{X}}/H \rightarrow \hat{\mathcal{X}}/G$.

It remains to find a cover $C_1 \rightarrow D_1$ as above. Consider the cover¹ defined by the extension $k(D_1) = k(u) \rightarrow k(C_1) = k(u, v)$, with $v^{p^2+p} + v = u$, where $p = \text{char}(k) > 0$. Then $C_1 \rightarrow D_1$ is étale outside of the pole y_1 of u . Let us show that the Galois closure $k(E)$ has positive genus. Let $t \in k(E)$ be such that $t^{p^2+p} + t = u$ and $t \neq v$. Let $w = t - v$. Then w satisfies the equation $(w^{p+1} - wt^p - w^p t)^p - w = 0$.

¹This example is given by Michel Matignon.

Hence $w = z^p$ where $z = w^{p+1} - wt^p - w^p t$. We have $z = z^{p^2+p} - z^p t^p - z^{p^2} t$, so

$$(t/z^p)^p + (t/z^p) = 1 - (1/z)^{p^2+p-1}.$$

This equation defines a p -cyclic cover $E' \rightarrow \mathbb{P}_k^1$, with conductor $m = p^2 + p - 1$ at $z = 0$, and étale elsewhere. Hence $g(E) \geq g(E') = (p-1)(m-1)/2 \geq 2$.

Remark 4.19 If we restrict ourself to the category of regular models of X , then Theorem 2.3 still holds. More precisely, given any model $\mathcal{X}$, there exists (after finite separable extension of K) a unique semi-stable and regular model dominating $\mathcal{X}$ and minimal for this property. This model is just the minimal desingularization of the stable hull of $\mathcal{X}$. However it does not commute with base change except when the normalization of $\mathcal{X}$ is smooth. On the other hand, Theorem 4.5 is no longer true in the setting of regular models. In general, given a morphism of models $\mathcal{X} \rightarrow \mathcal{Y}$, there is no finite morphisms of regular models dominating $\mathcal{X} \rightarrow \mathcal{Y}$, even after any finite extension of K (see [15], 6.5).

REFERENCES

- [1] S. Bosch, W. Lütkebohmert, *Stable reduction and uniformization of abelian varieties*, I, Math. Ann. **270** (1985), 349–379.
- [2] S. Bosch, W. Lütkebohmert, and M. Raynaud, *Néron Models*, Ergebnisse der Math., 3. Folge, **21**, Springer-Verlag 1990.
- [3] S. Bosch, W. Lütkebohmert, and M. Raynaud, *Formal and rigid geometry, IV, The reduced fibre theorem*, Invent. Math. **119** (1995), 361–398.
- [4] R. Coleman, *Stable maps of curves*, Documenta Math., extra volume K. Kato (2003), 217–225.
- [5] B. Conrad, B. Edixhoven and W. Stein, *$J_1(p)$ has connected fibers*, Documenta Math. **8** (2003), 331–408.
- [6] Ch. Deninger, A. Werner, *Vector bundles on p -adic curves and parallel transport*, preprint (2004), arxiv.org/abs/math/0403516.
- [7] H.P. Epp, *Eliminating Wild Ramification*, Invent. Math., **19** (1973), 235–249.
- [8] H. Grauert, R. Remmert, *Über die Methode der diskret bewerteten Ringe in der nicht-archimedischen Analysis*, Invent. Math. **2** (1966), 87–133.
- [9] Y. Henrio, *Arbre de Hurwitz et automorphismes d'ordre p des disques et des couronnes p -adiques formels*, thèse Université Bordeaux 1 (1999).
- [10] A.J. de Jong, *Smoothness, semi-stability and alterations*, Publ. Math. IHES, **83** (1996), 51–93.
- [11] A.J. de Jong, *Families of curves and alterations*, Ann. Inst. Fourier **47** (1997), 599–621.
- [12] F.-V. Kuhlmann, *A correction to: "Elimination of wild ramification" by H. P. Epp*, Invent. Math., **153** (2003), 679–681.
- [13] Q. Liu, *Algebraic geometry and arithmetic curves*, Oxford Grad. Texts in Math., **6**, Oxford Univ. Press, 2002.
- [14] Q. Liu, *Reduction and lifting of finite covers of curves*, Proceedings of the 2003 Workshop on Cryptography and Related Mathematics, Chuo University (2003), 161–180.
- [15] Q. Liu, D. Lorenzini, *Models of curves and finite covers*, Compos. Math. **118** (1999), 61–102.
- [16] S. Mochizuki, *The geometry of the compactification of the Hurwitz scheme*, Publ. Res. Inst. Math. Sci., Kyoto University, **31** (1995), 355–441.
- [17] M. Raynaud, *p -groupes et réduction semi-stable des courbes*, The Grothendieck Festschrift, vol. III (P. Cartier and al, eds), Birkhäuser, Prog. in Math. **88** (1990), 179–197.
- [18] J.-P. Serre, *Corps locaux*, Hermann, Paris, 1968.

CNRS, LABORATOIRE A2X, INSTITUT DE MATHÉMATIQUES DE BORDEAUX, UNIVERSITÉ DE BORDEAUX I, 351, COURS DE LA LIBÉRATION, 33405 TALENCE, FRANCE